PART F. CHAPTER 3. BRAIN HEALTH

Table of Contents

Introduction F3-1

Review of the Science F3-3

Overview of Questions Addressed F3-3

Data Sources and Process Used to Answer Questions F3-4

Y µ • š] } What is the relationship between physical activity and cognition? F3-4

Question 2. depressed mood and depression? F3-25

Question 4 X What is the relationship between physical activity and sleep? F3-37

Needs for Future Research F3-46

References

thoughts and actions, and achieve academic goals, including school readiness and achievement. In late adulthood, we aim to avoid dementia and cognitive impairment. Across the lifespan, we strive to ensure high-quality brain health, as manifested by optimally functioning cognition, low levels of anxiety and feelings of depression, a positive assessment of perceptions of quality of life, and comfortable and effective sleep patterns. Despite these common goals, and the fact that recent research has provided

4. What is the relationship between physical activity and sleep?
    a) Is there a dose-response relationship for either acute bouts of physical activity, or regular physical activity? If yes, what is the shape of the relationship?
    b) Does the relationship vary by age, sex, race/ethnicity, socioeconomic status, or weight status?
    c) Does the relationship vary for individuals with normal to impaired sleep behaviors? If yes, for which sleep disorders?

Data Sources and Process Used to Answer Questions

The Brain Health Subcommittee determined that systematic reviews, meta-analyses, pooled analyses, and reports provided sufficient literature to answer all four research questions. The databases searched included P dtd,dCtoanando CINo

Strong evidence demonstrates that greater amounts of physical activity are associated with a reduced risk of developing cognitive impairment, including Alzheimer's disease. PAGAC Grade: Strong.

Moderate evidence indicates that moderate-to-vigorous physical activity interventions can improve cognition in individuals with dementia. PAGAC Grade: Moderate

Moderate evidence indicates that moderate-to-vigorous physical activity can have beneficial effects on cognition in individuals with diseases or disorders that impair cognitive function, including attention deficit hyperactivity disorder, schizophrenia, multiple sclerosis, Parkinson's disease, and stroke. However,, mu[illegible]

cogn have annon(t)-2.998.(an)4.00e.8 (,)10.996au (iti)12.00sm (o)- (a,)9.005 (can)2.99cer)e.

anncodc 7oannitiouie.

term that represents a number of goal-directed processes that support thinking, reasoning, and problem solving), and dementia diagnoses. However, cognition as defined in this question does not include measurement of intelligence, motor function, personality, mood (addressed below in Question 3), and sensory and perceptual function.

In young and mid-life adults, effect sizes ranged from 0.12 to 0.15[4,5] physical activity improving cognition. Effects were largest for the cognitive domains of executive function, attention, processing speed,[5] and short-term memory.[4] In cognitively normal older adults, effect sizes ranged from non-significant[7] to .20[8] to 0.48[6] in favor of physical activity interventions positively influencing cognitive outcomes. Effect sizes were greatest for measures of executive function,[6] global cognition,[8] and attention.[7]

Impaired cognitive function: Strong evidence demonstrates that greater amounts of physical activity and lob 11.0eb2

Dose-response. The included reviews rarely report whether a dose-response relationship was observed for the effects of physical activity on cognitive outcomes. However, one meta-analysis[6] suggested that among older adults, larger effects on cognition were observed in randomized controlled trials in which physical activity bouts lasted 46-60 minutes in duration (compared to bout durations lasting 15-30 minutes and 31-45 minutes) and the interventions occurred for at least 6 months compared to interventions lasting 1-3 and 4-6 months. In addition, physical activity has a general effect across the aspects of cognition that were studied (i.e., executive, controlled, spatial, and speed), but the effect was selectively and disproportionately larger for tasks requiring greater amounts of executive control.[6]

Acute bouts of physical activity. Studies demonstrate a small, transient improvement in cognition following the cessation of a single, acute bout of physical activity, with effect sizes ranging from 0.014 to 0.67.[26-29] Reported effects were most consistent for domains of executive function,[26-29] but significant benefits were also realized for processing speed, attention, memory, and crystalized intelligence, the latter of which is a measure of general and verbal knowledge (e.g., what is the name of the first president of the United States).[26, 27, 29] Larger effects were also realized for preadolescent children and older adults relative to adolescents and young adults.[29]

Exercise intensity of an acute bout of activity had an effect on changes in cognition, with some findings suggesting an inverted-U shaped curve, as moderate-intensity exercise demonstrated a larger effect than light- and vigorous-intensity exercise,[27-29] and other studies indicating that very light-, light-, and moderate-intensity exercise benefited cognition, but hard-, very hard-, and maximal-intensity exercise intensity demonstrated no benefit.[26] The timing of the assessment of cognition relative to the cessation of the acute bout of exercise also demonstrated differences in the magnitude of the effect, with negative effects in cognition observed during the first 10 minutes following the exercise bout and the largest positive effect observed from 11 to 20 minutes and a smaller effect observed after 20 minutes following the acute phyt58n >>BDCnd a ued ce8.995 (th)7. T* (-)1.649 (in)t3f4e (t)6.m2.998 (in) (t)-2.9983 1

inadequate blinding, and variability in control group conditions. As such, the studies included in these meta-analyses and systematic reviews generally have a high risk of bias and low precision. However, despite these limitations, these studies appear to have high applicability, generalizability, and consistency. The effects are also detectable using acute exercise paradigms, where preadolescent children and older adults demonstrate large and consistent positive effects of moderate-intensity physical activity,[26-29] with some evidence to support 11 to 20 minutes in duration as being optimal for cognitive outcomes.[26]

For additional details on this body of evidence, visit https://health.gov/paguidelines/second-edition/report/supplementary-material.aspx for the Evidence Portfolio.

Comparing 2018 Findings with the 2008 Scientific Report

The 2008 Scientific Report[1] concluded that strong evidence demonstrated that physical activity delays the incidence of dementia and the onset of cognitive decline associated with aging. It also indicated that physical activity improves cognitive symptoms associated with dementia. Thus, the evidence described here considerably expands that described in 2008 by including significantly more observational studies and RCTs. This research finds that physical activity influences cognitive function across the lifespan, including both cognitively normal and impaired populations (e.g., schizophrenia). The effects are consistent across a variety of methods for assessing cognition (e.g., academic achievement and dementia diagnoses). The 2018 Scientific Report also demonstrates, for the first time, the positive effects of physical activity on biomarkers of brain health obtained from neuroimaging techniques (e.g., brain volume). Finally, the 2018 Scientific Report describes evidence on acute bouts of activity for improving cognitive function.

Public Health Impact

In 2017,

physical inactivity in the United States could potentially prevent 230,000 cases. The results from the 2018 Scientific Report provide support for the argument that physical activity reduces the risk of Alzheimer's disease and other dementias and that increasing physical activity in individuals with Alzheimer's disease could improve cognitive function.

The public health impact of the results summarized in the 2018 Scientific Report goes beyond Alzheimer's disease and dementia by demonstrating that physical activity influences cognitive function in children and healthy older adults. For example, academic achievement is a predictor of future job opportunities[37] and adult health outcomes.[38,39] Thus, these findings, which indicate that increasing physical activity during childhood may positively influence cognition and academic achievement, may have further downstream effects on many features of adult health and quality of life

Healthy older adults, even in the absence of a dementia, often show evidence for cognitive losses and decline, especially on measures of processing speed, memory, and executive function.[40] It is estimated that by the year 2050, the population of adults older than 65 years in the United States will reach 83.7

Maintaining or improving QoL is a universal goal. Being physically active has been suggested as one way to enhance perceptions of, and feelings of, QoL. This question focuses on the scientific literature that describes QoL as experienced by the general population across the lifespan. It also includes an assessment of the effects of physical activity on QoL in individuals with mental health issues. QoL among individuals who have a chronic physical condition, such as diabetes or osteoarthritis, is considered in Part F. Chapter 10. Individuals with Chronic Conditions.

The literature reviewed here focused on QoL and HRQoL, specifically. Searches were not conducted on "well-being" or its derivatives, such as subjective well-being, positive well-being, or psychological well-being. Those well-being concepts typically blend cognitive/evaluative and affective components and this question is limited to the cognitive/evaluative aspects widely known as quality of life.

Literature Reviewed

To answer this question, the Subcommittee reviewed 18 systematic reviews across the following populations: older adults (general),[49-51] older adults (with dementia),[52] adults,[53-58] youth,[59] individuals with schizophrenia,[60,61] individuals with depression,[52] and 14 meta-analyses across the following populations: adults,[62-64] older adults (general),[65-70] older adults (with dementia),[71,72] adn MCID 28 >>BDC 0 0 1 rg /T3

The reviews covered the following timeframe: inception to July 2011,[69] July 2011 to October 2014,[64] and inception to 2015.[73]

QoL was most often conceptualized as HRQoL, and assessed with the SF-36, the SF-12, the WHOQoL-Bref, and the EuroQoL Group 5- Dimension Self-Report Questionnaire.[73]

Individuals with Depression

The reviews covered the following timeframes: inception to June 2013,[52] inception to May 2013,[74] and inception to January 2015.[75] Two of these reviews included 7 studies, and another that examined the effects of yoga included 12 RCTs.

The number of studies reviewed that included both physical activity and QoL (N=10 studies) is much smaller than the total number of studies in the systematic reviews, ranging from one study[75] that included only the Mental Component of the SF-36, four studies in older adults with depression,[52] and a meta-analysis including four studies comparing physical activity to non-active controls, one study comparing physical activity to antidepressant medication, and one with comparison to cognitive therapy for depression.[74]

QoL was conceptualized as HRQoL, and assessed in most cases with the SF-36.[52]

type of activity.[53, 57] Of the studies reviewed, only one presented specific information on the frequency, intensity, time, and type (FITT) principles for exercise prescription;[70] however, the FITT principles were not reported in relation to QoL outcomes.

Evidence on the Overall Relationship

General Population - Older Adults

Overall, results showed that physical activity consistently resulted in improvements in QoL in older adults. One meta-analysis reported that collectively, exercise programs (1,317 participants) improved the QoL (overall and health-related combined) of older adult participants (Z = 2.03), and the pooled standardized mean difference (SMD) was 0.86 (95% confidence interval (CI): 0.11-1.62).[68] In another meta-analysis, statistically significant improvements were found for the physical function subscale of tunct a00 (epo)-2.ea00 (epo)-8 (ere)-4.004 iepo2.998 (e)9.in relatio (e)9.00icallfat68r8.99.hg /Hcipc

.9g71688ic =007 (P)-0.998 (b-2.93 .11 HR:.9987410 (62.91)-29.35 (:sig5<<o.(ro5%ere)-4ic CI)11.73 (:.99873 (04 Tf 0)-60.3

Physical activity as part of other activities that involve mental and physical components, such as qigong and tai chi, hold great potential for improving QoL in both healthy and chronically ill individuals.[51] However, effect sizes were not included, it was not reported which of the subdomains of QoL were improved, and results and conclusions were not separated by healthy and chronically ill participants. Moreover, given the mind-body nature of these modes of physical activity, it is not clear whether changes in QoL would be the result of changes in physical activity or other components of the activity (e.g., breathing, meditation).

In a pooled analysis,[76] participants who were active for more than 150 minutes per week of physical activity but then dropped to fewer than 150 minutes per week from baseline to 6 months showed a 11.8 point drop ($P<0.001$) in SF-36 physical function scores. In contrast, those who were active for fewer than 150 minutes per week of physical activity but then increased to more than 150 minutes per week from baseline to 6 months showed an increase of 5.1 points in SF-36 physical function scores.[76] These results indicate the importance of maintaining physical activity for maintaining HRQoL in late adulthood.

The effects of physical activity on non-HRQoL domains are more equivocal. Studies examining non-HRQoL domains show consistent and positive associations between physical activity and the domains of functional capacity, general QoL, and autonomy. These domains have been related to QoL in the elderly However, few studies were methodologically rigorous. Effect sizes were generally small or moderate and varied widely between studies and across QoL domains.[43]

Among frail older adults, one review found no significant differences in QoL among studies that used water exercises, flexibility exercises, tai chi, and resistance exercises,[65] and others had too few studies to make a conclusion.[46-48] These studies of QoL were not intended to capture objective measures of physical function (e.g., balance, gait speed), as the measures of QoL were developed to assess perceptions of functioning. Thus, in the context of frail older adults, beneficial effects of physical activity on measures of physical function may not be immediately apparent on perceptions of functioning that are captured by common instruments assessing QoL.

In summary, the evidence points to a positive effect of physical activity on both overall and health-related QoL in older adults. Physical health-related QoL has been investigated more consistently than mental health-related QoL. The limited available literature suggests that the physical activity effects on physical and mental health composite scores appear to be similar in both direction and magnitude. There were insufficient studies and sample sizes to adequately analyze effects of different exercise

The other six reviews reported similarly positive associations between greater amounts of physical activity and higher assessments of QoL.[54-56, 58, 63, 64]

General Population - Youth

There was no evidence available on the relationship between physical activity and QoL among youth.

QoL in most reports.[52] One RCT comparing yoga to a relaxation control group showed improvement of 50 percent or greater on the mental QoL domain.[75]

Collectively, these studies provide limited evidence for a moderate effect size of physical activity on physical and mental domains, but not overall QoL outcomes for adults with depression, when compared to placebo or inactive controls. In older adults with clinical depression, limited evidence from a small number of controlled studies suggests that physical activity is associated with improved QoL outcomes.[52] Thus, advancing age may serve as a response modifier for the effects of physical activity on QoL, consistent with our report that physical activity has a strong positive effect on HRQoL in the non-depressed older population.

Evidence on Specific Factors

Dose-response: Meta-analyses did not report on the effect of different doses of physical activity on QoL outcomes.

Non-experimental evidence from ecological momentary assessments provides insight into relations between physical activity and subsequent affective responses over a 3-hour interval.[84] In 8 out of 11 studies, physical activity was associated with more pleasant and activated subsequent affective states following the activity bout. Results were mixed with regard to physical activity and unpleasant feelings. Two studies found no association between physical activity and subsequent unpleasant feelings and two studies found that physical activity was associated with reduced unpleasant feelings. A fifth study found that physical activity did not lead to acute reductions in unpleasant feelings but that people who were typically more active reported fewer unpleasant feelings in general.

Dose-response: Strong evidence shows an effect of physical activity on immediate affective responses and that this effect is moderated by the imposed dose of activity.

Evidence on Specific Factors

Demographic factors and other moderators: Little is known about the persistence of the effects of physical activity on affective states across time or how they might be moderated by individual variability in demographic or other biological or environmental factors.

Biomarkers: Insufficient evidence was available from the reviewed literature to determine whether physical activity modifies biomarkers of brain structure and function in the context of affect. There were no studies reviewed that examined brain measures or other biomarkers.

Anxiety

Here, the Subcommittee defines anxiety as a noticeable, psychophysiological emotional state, which is most often characterized by feelings of apprehension, fear or expectations of fear, worry, nervousness, and physical sensations arising from activation of the autonomic nervous system (e.g., increased muscle tension, elevated heart rate, sweating). This normal human emotion becomes pathological (i.e., clinical

majority of this work has examined the effects of exercise in individuals without elevated symptoms of anxiety and/or not diagnosed with any clinical anxiety disorders.

Evidence on the Overall Relationship

To examine the effects of acute exercise bouts on measures of state anxiety, the Subcommittee reviewed evidence from a meta-analysis of 36 RCTs (involving 1,233 individuals [726 females] examining the effects of acute exercise on state anxiety published since 1980.[86] Samples varied from

to 6 months, with a range of 1 to 7 training sessions per week. Outcome measures varied considerably, from assessments of anxiety symptoms to clinical assessments of anxiety; all were used to assess anxiety before and after the exercise (or control) interventions. Control comparisons involved standard care (most often pharmacotherapy or cognitive behavioral therapy), a waitlist group that is tested several times before beginning the intervention, a placebo group, or another exercise intervention.

Physical activity had a significant effect on the reduction of trait anxiety. One review[94] reported a moderate effect (Cohen d (d)=0.31 for non-RCT studies; d=0.45 from RCTs) and another review[99] reported a small-to-moderate effect for resistance exercise training (d=0.20). Reviews comparing the effects of exercise to other treatments[88, 89, 93, 94] consistently reported that exercise interventions were at least as effective as standard care treatment for anxiety and sometimes even better.[94] To use one example, a meta-analysis[88] of exercise compared to various control groups (including active treatments) on trait anxiety in patient populations showed that exercise was as efficacious as, and not inferior to,

the evidence indicates that exercise may have beneficial effects on PTSD symptoms and that regular physical activity may reduce risk of developing PTSD. The evidence also suggests that yoga may be useful (d=0.48) in alleviating PTSD symptoms, but the studies show little consistency regarding the type of yoga and the length of treatment.

Dose-response. Limited evidence suggests a dose-response effect of physical activity on either state or trait anxiety symptoms.

Evidence on Specific Factors

Demographic factors. Moderate evidence indicates that state anxiety reduction is moderated by sex and age such that females and those older than age 25 years show greater reductions in state anxiety after participating in physical activity.[86] Insufficient evidence was available from the examined literature

unique studies, with 2 studies included in both reviews[54,127,128]; all studies were experimental in design. Each of the systematic reviews included six longitudinal studies[129,130]. The review of reviews[96] included four systematic reviews that had appropriate exposures and outcomes for this question; the sum of RCTs included in each of the 4 reviews totaled 93. In all of the reviews, parameters of physical activity were obtained from a variety of self-report instruments. Similarly, symptoms of depression were assessed with a wide variety of tools, standard and non-standard.[131]

All five studies reported statistically significant reductions in depressive symptoms in the more physically active groups. One meta-analysis reported a Hedges' g=0.26 (95% CI: -0.43 to -0.08)[54,128]; another a standardized mean difference of -0.61 (95% CI: -1.06 to -0.16).[127] The review of reviews reported a statistically significant reduction in the standardized mean difference among the more physically active

this effect modification, there are other reports showing similar effects across males and females.[94] In any case, potential sex differences (or lack thereof) should be interpreted with caution because of the higher prevalence of depression and depressive symptoms in females. In contrast, little to no

2018 Scientific Report includes an assessment of acute bouts of physical activity on measures of affect and state anxiety. Finally, the 2018 Scientific Report also provides an examination of physical activity on reducing depression and state and trait anxiety across multiple age groups and populations (e.g., youth).

Public Health Impact

In the United States, fewer than half of children and adults engage in regular physical activity.[134]

activity, is needed. The results reported in this chapter clearly indicate that physical activity is an effective and robust approach for reducing the risk of depression that would clearly have downstream consequences for quality of life, health care costs, and job productivity. Furthermore, these results also demonstrate that physical activity is an effective approach for improving both anxiety and depressive symptoms (symptoms that often co-occur), with effect sizes that are similar to that of the most effective pharmaceutical approaches.

In sum, physical activity holds great promise as a means for preventing and treating common mood disorders that are a significant source of disability, lower quality of life, and increased health care burden.

Question 4: What is the relationship between physical activity and sleep?

a) Is there a dose-response relationship for either acute bouts of physical activity, or regular physical activity? If yes, what is the shape of the relationship?
b) Does the relationship vary by age, sex, race/ethnicity, socioeconomic status, or weight status?
c) Does the relationship exist for individuals with impaired sleep behaviors or disorders? If yes, for which sleep disorders?

Sources of evidence: Systematic reviews, meta-analyses

Conclusion Statements

Strong evidence demonstrates that both acute bouts of physical activity and regular physical activity improve sleep outcomes in adults. PAGAC Grade: Strong.

Moderate evidence indicates that longer duration acute bouts of physical activity and regular physical activity improve sleep outcomes. These positive effects are independent of exercise intensity. PAGAC

Moderate evidence indic
preserved across age an

awakenings. In contrast, physical activity bouts performed 3 to 8 hours before bedtime were associated with reduced REM sleep.[142]

Dose-response. Moderate evidence indicates a dose-response relationship between the length in minutes but not the intensity or modality of moderate-to-vigorous physical activity and sleep outcomes. In adults, this evidence is supported by analyses from 59 controlled studies (N=2,863 participants) in which the length in minutes of acute physical activity bouts was found to moderate the beneficial effects on sleep onset latency (less), total sleep time (more), slow wave sleep (more), and rapid eye movement sleep (less).[142] In terms of regular physical activity, limited but concordant evidence suggests that more minutes of moderate-to-vigorous physical activity in each individual session is also associated with greater beneficial effects on reducing sleep onset latency. Taken together, these findings provide consistent evidence for a relationship between greater length in minutes of moderate-to-vigorous physical activity bouts associated with benefits to multiple objective and physiological sleep outcomes. In contrast to the length of each physical activity session, the number of weeks of the exercise intervention had a small but statistically significant effect on total sleep time, but no effect on sleep quality, latency, or efficiency.[142]

Regular physical activity levels influence the response to an acute bout of physical activity on slow wave sleep. Among individuals with high baseline physical activity, ach d5 (tal)-2.998 (statv03.036 0t92.998 (p)2wi]T

Evidence on Specific Factors

Age: In adults, moderate evidence indicates that relationships between physical activity and sleep outcomes are consistent in their effects across young, middle-aged, and older men and women.[142, 155, 158, 162, 163] Consistent evidence indicates a reduced beneficial effect of greater physical activity amount on sleep latency with aging, consisting of a 0.15 standard deviation decrease in the beneficial effects of regular physical activity for every decile increase in mean age.[142] In contrast, age does not moderate the relationship between greater amounts of regular physical activity and its beneficial effects on total sleep time, sleep efficiency, and sleep quality.

In contrast to systematic reviews in adults that include many controlled intervention studies, in children and adolescents, studies examining the relationship between physical activity and sleep are mostly cross-sectional, with a few cohort studies.[152, 155, 159] A meta-analysis of 15 studies of 12,604 individuals ages 14 to 24 years, reported a beneficial effect of physical activity on sleep with an overall standard mean difference of 0.77 (95% CI: 0.41-1.13).[155] Another meta-analysis of 11 cross-sectional studies reported a relationship betwen9rg/(e)-2.mf-3.995 (5 (9)7.002f)9.004cdacti6 98 (4)-3.0actiT04 (1)-2.9/-.ned a C

155

daytime sleepiness, compared to controls.[154] Another meta-analysis of 180 participants in 6 RCTs and 2 pre-post studies (the pre-post studies contributed 10 percent of the total number of participants) reported a decrease in AHI (unstandardized mean difference (USMD) =-0.536 (95% CI: -0.865 to -0.206) and reduced Epworth sleepiness scale (USMD=-1.246; 95% CI: -2.397 to -0.0953).[151] Finally, a network meta-analysis compared the effectiveness of supervised aerobic exercise training with continuous positive airway pressure (CPAP), mandibular advancement devices (MAD), and weight loss on AHI.[153]

Comparing 2018 Findings with the 2008 Scientific Report

The 2008 Scientific Report[1] concluded that "A small number of observational, population-based studies provides initial evidence supporting a positive association of regular participation in physical activity with lower odds of disrupted or insufficient sleep, including sleep apnea." The 2008 Scientific Report[1] also concluded that "a small number of RCTs supports the conclusion that regular participation in physical activity has favorable effects on sleep quality and is a useful component of good sleep hygiene."

environmental risk factors (e.g., stroke risk factors, traumatic brain injury) as covariates influencing the response to physical activity. To date, although candidate biomarkers and environmental risk factors have been identified, little systematic study in humans has emerged in the literature especially in relation to markers associated with affect, anxiety, depression, and sleep.

5. Conduct studies to monitor sedentary time and conduct randomized controlled trials that systematically reduce sedentary behaviors to improve the understanding of the impact of varying contexts, patterns, and durations of sedentary behavior on brain health outcomes (e.g., depression symptoms) throughout the lifespan and in populations with brain health disorders and diseases.

   Rationale: The understanding of the effects of sedentary behavior on brain health is in its infancy. Given that recent evidence indicates that sedentary behavior is distinct from physical inactivity, a greater understanding of the effect of sedentary behavior on brain health may inform and target interventions aimed at improving brain health across a variety of populations, including school-aged children, middle-aged adults, and older adults, as these populations spend considerable time during their day engaged in sitting and other sedentary behaviors. In addition, portable health technologies that continuously measure physical activity, estimate its intensity, and characterize sleep behavior, may offer inroads to better understand such relationships, and perhaps test novel interventions using connected health approaches.

6. Conduct appropriate analyses to examine effect modification by demographic factors. Such analytical approaches require studies that include large samples and substantial variation in sample characteristics (i.e., race/ethnicity, socioeconomic status).

   Rationale: Although some understanding of the effects of physical activity during the developing years and in aging has emerged, evidence for other demographic factors has not been demonstrated in a systematic fashion, affording little opportunity to form strong conclusions about

periods, and a broader range of instruments and outcomes relevant for brain health (e.g., mental subdomain of health-related quality of life, affect).

Rationale: To date, the temporal dynamics of the effects of physical activity on brain health are poorly understood. Yet, it is known that individuals start and stop exercise regimens on a regular basis and such variability in the consistency of physical activity may differentially influence the impact of physical activity on brain health outcomes. It is possible that the persistence of the effects might also depend on the dose of activity (frequency, intensity, time, type), the age of the individual, the presence of a disorder or disease, or other factors. Enrolling samples of sufficient size to support mediator analyses (i.e., exploration of putative mechanisms through which the interventions operate) will provide useful information for adapting the interventions to optimize uptake among different subgroups as well as to identify key elements that are essential to improving brain health.

8. Conduct randomized controlled trials and prospective observational research on the impact of muscle-strengthening exercises (often referred to in the literature as resistance training) and other

4. Roig M, Nordbrandt S, Geertsen SS, Nielsen JB. The effects of cardiovascular exercise on human memo

17. Tan BWZ, Pooley JA, Speelman CP. A meta-analytic review of the efficacy of physical exercise interventions on cognition in individuals with autism spectrum disorder and ADHD. J Autism Dev Disord 2016;46(9):3126-3143. doi:10.1007/s10803-016-2854-

(iso)12.0hc)-2.18 (y)-3.8 Beckett MW, Ardern CI, Rotondi M2.998 (i)7l(is)12.002 (i18 (ern CI)12.518 (s). at13.33 (-)12.519 (an)4.004

31. Li MY, Huang MM, Li SZ, Tao J, Zheng GH, Chen LD. The effects of aerobic exercise on the structure and function of DMN-related brain regions: A systematic review. Int J Neurosci. 2016;127(7):634-649.

32. Dinoff A, Herrmann N, Swardfager W, et al. The effect of exercise training on resting concentrations of peripheral brain-derived neurotrophic factor (BDNF): A meta-analysis. PLoS One 2016;11(9):e0163037. doi: https://doi.org/10.1371/journal.pone.0163037

33. Halloway S, Wilbur J, Schoeny ME, Arfanakis K. Effects of endurance-focused physical activity interventions on brain health: A systematic review. Biol Res Nurs

46. Clegg AP, Barber SE, Young JB, Forster A, Iliffe SJ. Do home-based exercise interventions improve outcomes for frail older people? Findings from a systematic review. Rev Clin Gerontol. 2012;22(1):68-78.

47. Chao YY, Scherer YK, Montgomery CA. Effects of using Nintendo Wii exergames in older adults: a review of the literature. J Aging Health. 2015;27(3):379-402. doi:10.1177/0898264314551171.

48. Weening-Dijksterhuis E, de Greef MH, Scherder EJ, Slaets JP, van der Schans CP. Frail institutionalized older persons: a comprehensive review on physical exercise, physical fitness, activities of daily living, and quality-of-life. Am J Phys Med Rehabil. 2011;90(2):156-168. doi:10.1097/PHM.0b013e3181f703ef.

49. Stevens Z, Barlow C, Kendrick D, et al. Effectiveness of general practice-based physical activity promotion for older adults: systematic review (1)6.993 (0)-2.998 (.15 (P)-3.9.(1)6.9936 Td (of)98 :(in)5 (g)3m (a

old(rick D)-1 .00 (.1)9.004 (0)-2.998 (9)7.002 (3)-2.9986/0jang013.0bef.0 8.841EMC /Lbl <</MCID 19 >>BDC2

60. Vancampfort D, Probst M, Helvik Skjaerven L, et al. Systematic review of the benefits of physical therapy within a multidisciplinary care approach for people with schizophrenia. Phys Ther 2012;92(1):11-23. doi:10.2522/ptj.20110218.

61. Vera-Garcia E, Mayoral-Cleries F, Vancampfort D, Stubbs B, Cuesta-Vargas AI. A systematic review of the benefits of physical therapy within a multidisciplinary care approach for people with schizophrenia: An update. Psychiatry Res. 2015;229(3):828-839. doi:10.1016/j.psychres.2015.07.083.

62. Gillison FB, Skevington SM, Sato A, Standage M, Evangelidou S. The effects of exercise interventions on quality of life in clinical and healthy populations; a meta-analysis. Soc Sci Med. 2009;68(9):1700-1710. doi:10.1016/j.socscimed.2009.02.028.

63. Soga M, Gaston KJ, Yamaura Y. Gardening is beneficial for health: a meta-analysis. Prev Med Rep. 2017;5:92-99. doi:10.1016/j.pmedr.2016.11.007.

64. Wang F, Man JK, Lee EK, et al. The effects of Qigong on anxiety, depression, and psychological well-being: a systematic review and meta-analysis. Evid Based Complement Alternat Med. 2013;2013:152738. doi:10.1155/2013/152738.

65. Chou CH, Hwang CL, Wu YT. Effect of exercise on physical function, daily living activities, and quality of life in the frail older adults: a meta-analysis. Arch Phys Med Rehabil. 2012;93(2):237-244. doi:10.1016/j.apmr.2011.08.042.

66. Forsman AK, Nordmyr J, Wahlbeck K. Psychosocial interventions for the promotion of mental health and the prevention of depression among older adults. Health Promot Int. 2011;26(Suppl 1):i85-i107. doi:10.1093/heapro/dar074.

67. Kelley GA, Kelley KS, Hootman JM, Jones DL. Exercise and health-related quality of life in older community-dwelling adults: a meta-analysis of randomized controlled trials. J Appl Gerontol 2009;28(3):369-394. doi: https://doi.org/10.1177/0733464808327456.

68. Park SH, Han KS, Kang CB. Effects of exercise programs on depressive symptoms, quality of life, and self-esteem in older people: a systematic review of randomized controlled trials. Appl Nurs Res 2014;27(4):219-226. doi:10.1016/j.apnr.2014.01.004.

69. Patel NK, Newstead AH, Ferrer RL. The effects of yoga on physical functioning and health related quality of life in older adults: a systematic review and meta-analysis. J Altern Complement Med 2012;18(10):902-917. doi:10.1089/acm.2011.0473.

70. Raymond MJ, Bramley-Tzerefos RE, Jeffs KJ, Winter A, Holland AE. Systematic review of high-intensity progressive resistance strength training of the lower limb compared with other intensities of strength training in older adults. Arch Phys Med Rehabil. 2013;94(8):1458-1472. doi:10.1016/j.apmr.2013.02.022.

71. Potter R, Ellard D, Rees K, Thorogood M. A systematic review of the effects of physical activity on physical functioning, quality of life and depression in older people with dementia. Int J Geriatr Psychiatry 2011;26(10):1000-1011. doi:10.1002/gps.2641.

72. Ojagbemi A, Akin-Ojagbemi N. Exercise and quality of life in dementia. penti ahv(1 cise e(J)3.9o(1)9.004 (0) 3.007 s02 (m)6.005 (atic 2.998 (o)-7.1:)4.004 (5.007 (e)8.994 (sych2

73. Dauwan M, Begemann MJ, Heringa SM, Sommer IE. Exercise improves clinical symptoms, quality of life, global functioning, and depression in schizophrenia: a systematic review and meta-analysis. Schizophr Bull 2016;42(3):588-599. doi:10.1093/schbul/sbv164.

74. Cooney GM, Dwan K, Greig CA, et al. Exercise for depression. Cochrane Database Syst Rev 2013;(9):Cd004366. doi:10.1002/14651858.CD004366.pub6.

75. Cramer H, Lauche R, Langhorst J, Dobos G. Yoga for depression: a systematic review and meta-analysis. Depress Anxiety 2013;30(11):1068-1083. doi:10.1002/da.22166.

88. Bartley CA, Hay M, Bloch MH. Meta-analysis: aerobic exercise for the treatment of anxiety disorders. Prog Neuropsychopharmacol Biol Psychiatry. 2013;45(2):34-39. doi:10.1016/j.pnpbp.2013.04.016.

89. Stubbs B, Vancampfort D, Rosenbaum S, et al. An examination of the anxiolytic effects of exercise for people with anxiety and stress-related disorders: a meta-analysis. Psychiatry Res. 2017;49:102-108. doi:10.1016/j.psychres.2016.12.020.

90. Gordon BR, McDowell CP, Lyons M, Herring MP. The effects of resistance exercise training on anxiety: a meta-analysis and meta-regression analysis of randomized controlled trials. Sports Med. 2017;doi:10.1007/s40279-017-0769-

91. Mochcovitch MD, Deslandes AC, Freire RC, Garcia RF, Nardi AE. The effects of regular physical activity on anxiety symptoms in healthy older adults: a systematic review. Rev Bras Psiquiatr. 2016;38(3):255-261.

92. Jayakody K, Gunadasa S, Hosker C. Exercise for anxiety disorders: systematic review. Br J Sports Med. 2014;48(3):187-196. doi:10.1136/bjsports-2012-091287.

93. Stonerock GL, Hoffman BM, Smith PJ, Blumenthal JA. Exercise as treatment for anxiety: systematic review and analysis. Ann Behav Med. 2015;49(4):542-556. doi:10.1007/s12160-014-9685-9.

94. Wegner E09-man B11c.005 (tm)3.995 (J)0, Sx2.998 ((98 (er)7H125 >>BDC 11)11.005 (y)J)0, Sx (l J)14.

103. Robertson R, Robertson A, Jepson R, Maxwell M. Walking for depression or depressive symptoms: a systematic review and meta-analysis. Ment Health Phys Act. 2012;5(1):66-75.

104. de Souza Moura AM, Lamego MK, Paes F, et al. Effects of aerobic exercise on anxiety disorders: a systematic review. CNS Neurol Disord Drug Targets. 2015;14(9):1184-1193. doi:10.2174/1871527315666151111121259.

105. Nystrom MB, Neely G, Hassmen P, Carlbring P. Treating major depression with physical activity: a systematic overview with recommendations. Cogn Behav Ther. 2015;44(4):341-352. doi:10.1080/16506073.2015.1015440.

106. Rebar AL, Stanton R, Geard D, Short C, Duncan MJ, Vandelanotte C. A meta-meta-analysis of the effect of physical activity on depression and anxiety in non-clinical adult populations. Health Psychol Rev 2015;9(3):366-378. doi:10.1080/17437199.2015.1022901.

107. Yan S, Jin Y, Oh Y, Choi Y. Effect of exercise on depression in university students: a meta-analysis of randomized controlled trials. J Sports Med Phys Fitness. 2016;56(6):811-816.

108. Josefsson T, Lindwall M, Archer T. Physical exercise intervention in depressive disorders: meta-analysis and systematic review. Scand J Med Sci Sports. 2014;24(2):259-272. doi:10.1111/sms.12050.

109. Schuch FB, Vancampfort D, Richards J, Rosenbaum S, Ward PB, Stubbs B. Exercise as a treatment for depression: a meta-analysis adjusting for publication bias. J Psychiatr Res. 2016;77:42-51. doi:10.1016/j.jpsychires.2016.02.023.

110. Farah, WH, Alsawas, M, Mainou, M, et al. Non-pharmacological treatment of depression: a systematic review and evidence map. Evid Based Med. 2016;21(6):214-221.

111. Mura G, Carta MG. Physical activity in depressed elderly. A systematic review. Clin Pract Epidemiol Ment Health. 2013;9:125-135. doi:10.2174/1745017901309010125.

112. Schuch FB, Vancampfort D, Rosenbaum S, et al. Exercise for depression in older adults: a meta-analysis of randomized controlled trials adjusting for publication bias. Rev Bras Psiquiatr 2016c;38(3):247-254. doiw.

syid12.998 ((w)-4.004 T1 11.04 Tf 355.055 0 T1,CR)10 (ev J)8.995 (BAg)2.998 ((8 (gP)3.995 (r))14.004 y)-

Part F. Chapter 3.

145. American Academy of Sleep Medicine. International classification of sleep disorders, 3rd ed. Darien, IL: American Academy of Sleep Medicine; 2014.

146.